

THE CHROMOSOMES OF THE CHICK SOMA.

ROBERT T. HANCE,

THE ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH AND THE ZOÖLOGICAL
LABORATORY, UNIVERSITY OF PENNSYLVANIA.¹

The accumulated evidence in favor of the chromosomes being in some way concerned with the determination and control of the somatic characters has become so impressive as to be almost unassailable despite the possible objection of a few that the last link in the chain (a demonstration of the actual activity of these nuclear bodies as hereditary character bearers), due to the deficiencies in our knowledge of cell chemistry, has yet to be forged. This latter point is one about which geneticists and cytologists are, at present, not greatly concerned and conclusive data upon the physiological behavior of the chromosomes does not seem at all imminent. While information is gradually accumulating in preparation for this final analysis much may still be accomplished by purely morphological studies with tentative interpretations of physiological activity based on these observations. A recent study of the chromosomes associated with sex in the chick (5) involved extensive observations on the chromosomes of dividing somatic cells, the results of which are recorded below.

OBSERVATIONS.

The technique of preparation (3), of counting and measuring the chromosomes, the amount and source of the material (5) have all been recorded and need not be repeated here. The chromosomes of over 150 cells have been drawn and studied. As the nervous system is the most actively growing of all the regions of the body the majority of the cells studied have naturally been found in some part of it. Cells with clear polar views of metaphase chromosomes have been found in the brain, neural tube,

¹ The observations recorded in this paper were made principally at the University of Pennsylvania during the writer's tenure of a National Research Council Fellowship.

optic cup, auditory vesicle, connective tissue, heart muscle, blood, amnion, gonads and in tissue cultures of muscles.

The Chromosome Number.—The difficulties involved in determining the exact number of chromosomes in the chick have been described in detail (5). Briefly, the trouble encountered may be said to be due, first to the extreme smallness of the shortest chromosomes of the complex making observation at times uncertain and second, to the apparent failure of the component granules of the smaller chromosomes to unite or at least to clearly indicate their proper relations to each other until late metaphase if indeed it really and always happens then.

The average number of chromosomes in the soma of the chick, based on 78 counts, is 33. I think that this is perhaps lower than the actual number which the most satisfactory counts indicate to be about 35 or 36. This variation in the chick is apparently due to a failure of the chromomeres or parts of the smaller chromosomes to unite rather than to fragmentation as in the case of the pig (1). This opinion is based upon the observations in the chick of larger numbers of distinct chromatin bodies (with occasional visible connecting threads between them) in the prophase than could be found in cells in later stages of mitosis whereas in the pig a reduction in the chromosome number as the metaphase was neared did not occur. The smaller chromosomes of the chick are the only ones concerned while in the pig the long ones are the ones that become broken up.

The Chromosome Form.—All of the longer chromosomes (about 12 in number) of the somatic cells are in the form of J's while the shorter ones are rods. Their structure and size relations (5) are alike in all the tissues studied including tissue cultures (4). The largest chromosome (chromosome pair in the male) of the complex has been shown to be the one associated with sex and is found in all somatic cells as clearly as in the cells of the gonads (5), Figs. 1 to 9. The female embryonic cells are heterozygous for this chromosome while the male cells are homozygous. No differences in the chromosomes or their behavior have been noted between any of the body tissues or in comparison with those of the embryonic germ cells.

DISCUSSION.

The morphological data on the behavior of the chromosomes in the developing embryo, although admittedly scant, has so far given no clue to the manner in which the chromosomes may contribute their potentialities to the growing organism. Before a study of somatic chromosomes had been made it seemed reasonable to expect to find the various highly differentiated cells of the body with chromosome numbers, morphology or behavior at variance both with those found in other tissues and with the specific number and general characteristics found in the gonads. This has been found not to be the case in at least three forms, the pig (1), the evening primrose (2) and the chick (5). There was some fragmentation of the somatic chromosomes in the first two forms but it was shown that none of the chromatin was lost. Furthermore this fragmentation was neither specific for any particular tissue nor constant in amount. In general the chromosome situation in the soma seems to be entirely similar to that found in the unreduced gonad cells. This is a matter difficult to understand in the present state of our information. Of all the characteristics that must be controlled or borne by or at least associated with the chromosomes (as shown by the studies on *Drosophila*) few have any chance for expression in the majority of the somatic cells and tissues and must therefore be inhibited in one way or another. The cells of the lower forms of animal life largely retain the germ-like power of reproducing the portion lost or even an entire organism following injury. This power is perhaps even more marked in plants. As differentiation becomes more extreme in the animal kingdom the ability to regenerate a part or a whole animal from cells already specialized becomes less and less until in the highest types of animals somatic cells are usually able to produce only somatic cells like themselves. Yet the chromosomes in these highly specialized cells have, in the examples studied, been found to be entirely similar to those in the germ cells containing the possibilities for a complex animal or plant. This may suggest that the chromosomes are functionless in the differentiated Soma. Or it may be that having contributed their share in the production of the specialized tissue are thereafter inactive as far as are concerned the general somatic

attributes or determiners with which originally they must have been equipped. If the latter interpretation is correct the perpetuation of the complete mitotic mechanism in the soma as it exists in the reproductive organs may seem, as far as any need for an exact division of the genes is concerned, somewhat unnecessary, to be classed possibly with vestigial organs and having no more significance. In view of the great delicacy of the mechanism that exists in all cells for accurately dividing the chromatin the last suggestion does not seem impressive. But as far as our morphological data on the behavior of somatic chromosomes go, together with the behavior of the soma in growth and regeneration, the above suggestion is at least a possibility to be considered in our attempt to get at the physiology of development and genetics.

SUMMARY.

1. The chromosome number in the somatic cells of the chick is about 35 or 36.
2. No characteristic differences between the number, the sizes, the morphology or the behavior of the chromosomes in comparison either with each other or with the cells of the gonads have been noted.
3. In view of the entire similarity of the somatic and germinal mitotic behavior and in consideration of the complete inability of highly specialized cells to regenerate other than cells similar to themselves it is tentatively suggested as a basis for future discussion that the somatic chromosomes, as far at least as their genetic function is concerned, have either become functionless or their cytoplasmic environment is incapable of reacting to the possibilities presumably carried by them.

BIBLIOGRAPHY.

1. Hance, Robert T.
'17 The Diploid Chromosome Complexes of the Pig (*Sus scrofa*) and their Variations. *J. Morph.*, Vol. 30, p. 156-222.
2. Hance, Robert T.
'18 Variations in the Somatic Chromosomes of *Oenothera scintillans* DeVries. *Genetics*, Vol. 3, p. 225-275.
3. Hance, Robert T.
'25 The Fixation of Avian Chromosomes. *Anat. Rec.*, Vol. 31, p. 87-92.
4. Hance, Robert T.

- '26 A Comparison of Mitosis in Chick Tissue Cultures and in Sectioned Embryos. *BIOL. BULL.*, Vol. L., p. 155-159.
5. **Hance, Robert T.**
- '26 Sex and the Chromosomes in the Domestic Fowl (*Gallus domesticus*). *J. Morph.* (In press.)
6. **Lipschutz, Alexander.**
- '24 The Internal Secretions of the Sex Glands. Wilkins and Wilkins, Baltimore.

DESCRIPTION OF PLATE I.

All figures were drawn at table level with a Zeiss 1.5 mm. apochromat and a 15 \times Orthoscopic ocular. They are reproduced at the size originally drawn. All are polar views of metaphase plates in chick embryonic cells with the exception of Fig. 5.

FIG. 1. From optic cup. 35 chromosomes.

FIG. 2. From amnion. 30 chromosomes.

FIG. 3. From a male gonad. 39 chromosomes.

FIG. 4. From neural tube. 32 chromosomes.

FIG. 5. Late prophase in connective tissue cell grown in a tissue culture. 38 chromosomes.

FIG. 6. From heart muscle. 33 chromosomes.

FIG. 7. From brain. 35 chromosomes.

FIG. 8. From female gonad. 36 chromosomes.

FIG. 9. From neural tube. 35 chromosomes.



1



2



3



4




5



6



7



8



9

BIOLOGICAL BULLETIN

OF THE

Marine Biological Laboratory

WOODS HOLE, MASS.

VOL. LI

JULY, 1926

No. I

CONTENTS



Twenty-Eighth Annual Report of the Marine Biological Laboratory. I

PUBLISHED MONTHLY BY THE
MARINE BIOLOGICAL LABORATORY

PRINTED AND ISSUED BY
LANCASTER PRESS, INC.
LANCASTER, PA.

AGENT FOR GREAT BRITAIN
WHELDON & WESLEY, LIMITED
2, 3 and 4 Arthur Street, New Oxford Street, London, W. C. 2

Single Numbers, \$1.00. Per Volume (6 numbers), \$4.50

Editorial Staff

- GARY N. CALKINS—*Columbia University.*
E. G. CONKLIN—*Princeton University.*
M. H. JACOBS—*University of Pennsylvania.*
C. R. MOORE—*University of Chicago.*
GEORGE T. MOORE—*The Missouri Botanic Garden.*
T. H. MORGAN—*Columbia University.*
W. M. WHEELER—*Harvard University.*
E. B. WILSON—*Columbia University.*
-

Managing Editor

- FRANK R. LILLIE—*The University of Chicago.*
-

All communications and manuscripts should be sent to the Managing Editor, the University of Chicago, Sept. 15th to June 15th, or Woods Hole, Mass., June 15th to Sept. 15th. Subscriptions and other matter should be addressed to the Biological Bulletin, Prince and Lemon Streets, Lancaster, Pa.

BIOLOGICAL BULLETIN

OF THE

Marine Biological Laboratory

WOODS HOLE, MASS.

VOL. LI

AUGUST, 1926

No. 2

CONTENTS

CASTLE, EDWARD S.	<i>Observations on Motility in Certain Cyanophyceæ</i>	69
HARMAN, MARY T., AND ROOT, FRANK P.	<i>Number and Behavior of the Chromosomes in Cavia cobaya (the Common Guinea Pig)</i> ..	73
HARVEY, E. NEWTON.	<i>On the Inhibition of Animal Luminescence by Light</i>	85
HARVEY, E. NEWTON.	<i>Oxygen and Luminescence, with a Description of Methods for Removing Oxygen from Cells and Fluids</i>	89
ESSENBERG, J. M.	<i>Complete Sex-Reversal in the Viviparous Teleost Xiphophorus helleri</i>	98
MOORE, CARL R.	<i>Scrotal Replacement of Experimental Cryptorchid Testes and the Recovery of Spermatogenic Function (Guinea Pig)</i>	112
LAWRENCE, WALTER.	<i>The Fate of the Germinal Epithelium of Experimental Cryptorchid Testes of Guinea Pigs</i>	129

PUBLISHED MONTHLY BY THE
MARINE BIOLOGICAL LABORATORY

PRINTED AND ISSUED BY
LANCASTER PRESS, INC.
LANCASTER, PA.

AGENT FOR GREAT BRITAIN
WHELDON & WESLEY, LIMITED
2, 3 and 4 Arthur Street, New Oxford Street, London, W. C. 2

Single Numbers, \$1.00. Per Volume (6 numbers), \$4.50

Editorial Staff

GARY N. CALKINS—*Columbia University.*

E. G. CONKLIN—*Princeton University.*

M. H. JACOBS—*University of Pennsylvania.*

FRANK R. LILLIE—*University of Chicago.*

GEORGE T. MOORE—*The Missouri Botanic Garden.*

T. H. MORGAN—*Columbia University.*

W. M. WHEELER—*Harvard University.*

E. B. WILSON—*Columbia University.*

Managing Editor

C. R. MOORE—*The University of Chicago.*

All communications and manuscripts should be sent to the Managing Editor, the University of Chicago, Sept. 15th to June 15th, or to Biological Bulletin, Woods Hole, Mass., June 15th to Sept. 15th. Subscriptions and other matter should be addressed to the Biological Bulletin, Prince and Lemon Streets, Lancaster, Pa.

BIOLOGICAL BULLETIN

OF THE

Marine Biological Laboratory

WOODS HOLE, MASS.



VOL. LI

SEPTEMBER, 1926

No. 3

CONTENTS

- FROBISHER, MARTIN, JR. *Observations on the Relationship Between a Red Torula and a Mold Pathogenic for Drosophila melanogaster.* 153
- YOUNG, DONNELL B. *Nuclear Regeneration in Stylonychia mytilus.* 163
- MINNICH, DWIGHT ELMER. *The Chemical Sensitivity of the Tarsi of Certain Muscid Flies.* 166
- GREGORY, LOUISE H. *Effects of Changes in Medium During Different Periods in the Life History of Uroleptus mobilis.* 179
- YOUNG, WILLIAM C.,
AND PLOUGH, HAROLD H. *On the Sterilization of Drosophila by High Temperature* 189
- SKOWRON, STANISLAW. *On the Luminescence of Microscoplex phosphoreus Dug.* 199
- KATER, J. MCA. *Chromosomal Vesicles and the Structure of the Resting Nucleus in Phaseolus.* 209

PUBLISHED MONTHLY BY THE
MARINE BIOLOGICAL LABORATORY

PRINTED AND ISSUED BY
LANCASTER PRESS, INC.
LANCASTER, PA.

AGENT FOR GREAT BRITAIN
WHELDON & WESLEY, LIMITED
2, 3 and 4 Arthur Street, New Oxford Street, London, W. C. 2

Single Numbers, \$1.00. Per Volume (6 numbers), \$4.50

Editorial Staff

GARY N. CALKINS—*Columbia University.*

E. G. CONKLIN—*Princeton University.*

M. H. JACOBS—*University of Pennsylvania.*

FRANK R. LILLIE—*University of Chicago.*

GEORGE T. MOORE—*The Missouri Botanic Garden.*

T. H. MORGAN—*Columbia University.*

W. M. WHEELER—*Harvard University.*

E. B. WILSON—*Columbia University.*

Managing Editor

C. R. MOORE—*The University of Chicago.*

All communications and manuscripts should be sent to the Managing Editor, the University of Chicago, Sept. 15th to June 15th, or to Biological Bulletin, Woods Hole, Mass., June 15th to Sept. 15th. Subscriptions and other matter should be addressed to the Biological Bulletin, Prince and Lemon Streets, Lancaster, Pa.

BIOLOGICAL BULLETIN

OF THE

Marine Biological Laboratory

WOODS HOLE, MASS.

VOL. LI

OCTOBER, 1926

No. 4

CONTENTS

- LAMBERT, W. V.,
AND KNOX, C. W. *Genetic Studies in Poultry*..... 225
- McNALLY, EDNA. *Life Cycle of Nassula ornata and Nassula elegans: Are these Species Valid?*..... 237
- KUSNEZOV, N. J. *The Morphology of the Copulatory Structures in Some Cases of Gynandromorphism in Lepidoptera*..... 245
- BERRY, S. STILLMAN. *Note on the Occurrence and Habits of a Luminous Squid (Abralia veranyi) at Madeira*..... 257
- GABRITSCHESKY, E. *Convergence of Coloration Between American Pilose Flies and Bumblebees (Bombus)*..... 269

PUBLISHED MONTHLY BY THE
MARINE BIOLOGICAL LABORATORY

PRINTED AND ISSUED BY
LANCASTER PRESS, INC.
LANCASTER, PA.

AGENT FOR GREAT BRITAIN
WHELDON & WESLEY, LIMITED
2, 3 and 4 Arthur Street, New Oxford Street, London, W. C. 2

Single Numbers, \$1.00. Per Volume (6 numbers), \$4.50

Editorial Staff

GARY N. CALKINS—*Columbia University.*

E. G. CONKLIN—*Princeton University.*

M. H. JACOBS—*University of Pennsylvania.*

FRANK R. LILLIE—*University of Chicago.*

GEORGE T. MOORE—*The Missouri Botanic Garden.*

T. H. MORGAN—*Columbia University.*

W. M. WHEELER—*Harvard University.*

E. B. WILSON—*Columbia University.*

Managing Editor

C. R. MOORE—*The University of Chicago.*

All communications and manuscripts should be sent to the Managing Editor, the University of Chicago, Sept. 15th to June 15th, or to Biological Bulletin, Woods Hole, Mass., June 15th to Sept. 15th. Subscriptions and other matter should be addressed to the Biological Bulletin, Prince and Lemon Streets, Lancaster, Pa.

BIOLOGICAL BULLETIN

OF THE

Marine Biological Laboratory

WOODS HOLE, MASS.

VOL. LI

NOVEMBER, 1926

No. 5

CONTENTS

- BECKER, ELERY R. *The Flagellate Fauna of the Cæcum of the Striped Ground Squirrel, Citellus tridecemlineatus, with Special Reference to Chilomastix magna sp. nov.* 287
- BIGELOW, ROBERT PAYNE. *Variation in the Number of Fin-Rays in Three Species of Fundulus of the Woods Hole Region.* 299
- TANNREUTHER, GEORGE W. *Life History of Prorodon griseus.* 303
- BAKER, WOOLFORD B. *Studies in the Life History of Euglena.* 321
- BODINE, JOSEPH HALL. *Hydrogen Ion Concentration in the Blood of Certain Insects (Orthoptera)* 363
-

PUBLISHED MONTHLY BY THE
MARINE BIOLOGICAL LABORATORY

PRINTED AND ISSUED BY
LANCASTER PRESS, INC.
LANCASTER, PA.

AGENT FOR GREAT BRITAIN
WHELDON & WESLEY, LIMITED
2, 3 and 4 Arthur Street, New Oxford Street, London, W. C. 2

Single Numbers, \$1.00. Per Volume (6 numbers), \$4.50

Editorial Staff

GARY N. CALKINS—*Columbia University.*

E. G. CONKLIN—*Princeton University.*

M. H. JACOBS—*University of Pennsylvania.*

FRANK R. LILLIE—*University of Chicago.*

GEORGE T. MOORE—*The Missouri Botanic Garden.*

T. H. MORGAN—*Columbia University.*

W. M. WHEELER—*Harvard University.*

E. B. WILSON—*Columbia University.*

Managing Editor

C. R. MOORE—*The University of Chicago.*

All communications and manuscripts should be sent to the Managing Editor, the University of Chicago, Sept. 15th to June 15th, or to Biological Bulletin, Woods Hole, Mass., June 15th to Sept. 15th. Subscriptions and other matter should be addressed to the Biological Bulletin, Prince and Lemon Streets, Lancaster, Pa.



20712-56 117/27

BIOLOGICAL BULLETIN

OF THE

Marine Biological Laboratory

WOODS HOLE, MASS.

VOL. LI

DECEMBER, 1926

No. 6

CONTENTS

- WHITING, P. W. *Influence of Age of Mother on Appearance of an Hereditary Variation in Habrobracon*..... 371
- CALKINS, GARY N.,
AND BOWLING, RACHEL C. *Gametic Meiosis in Monocystis*..... 385
- GATES, G. E. *Preliminary Note on a New Protozoan Parasite of Earthworms of the Genus Eutyphæus* 400
- KINNEY, ELIZABETH. *A Cytological Study of Secretory Phenomena in the Silk Gland of Hyphantria cunea*..... 405
- HAMLETT, G. W. DELUZ. *The Linkage Disturbance Involved in the Chromosome Translocation I. of Drosophila, and its Probable Significance* 435
- HANCE, ROBERT T. *The Chromosomes of the Chick Soma*.. 443
-

PUBLISHED MONTHLY BY THE
MARINE BIOLOGICAL LABORATORY

PRINTED AND ISSUED BY
LANCASTER PRESS, INC.
LANCASTER, PA.

AGENT FOR GREAT BRITAIN
WHELDON & WESLEY, LIMITED
2, 3 and 4 Arthur Street, New Oxford Street, London, W. C. 2

Single Numbers, \$1.00. Per Volume (6 numbers), \$4.50

Editorial Staff

GARY N. CALKINS—*Columbia University.*

E. G. CONKLIN—*Princeton University.*

M. H. JACOBS—*University of Pennsylvania.*

FRANK R. LILLIE—*University of Chicago.*

GEORGE T. MOORE—*The Missouri Botanic Garden.*

T. H. MORGAN—*Columbia University.*

W. M. WHEELER—*Harvard University.*

E. B. WILSON—*Columbia University.*

Managing Editor


C. R. MOORE—*The University of Chicago.*

All communications and manuscripts should be sent to the Managing Editor, the University of Chicago, Sept. 15th to June 15th, or to Biological Bulletin, Woods Hole, Mass., June 15th to Sept. 15th. Subscriptions and other matter should be addressed to the Biological Bulletin, Prince and Lemon Streets, Lancaster, Pa.

Notice to contributors. Every paper to appear in the Biological Bulletin should be accompanied by an author's abstract presenting the chief results of the investigation. The abstract should not exceed 225 words in length.

For indexing purposes there should be, in addition to the title, one or more subject headings indicating in a word or two the divisions of the subject discussed in the paper. The entire name of the author (of each author if a joint paper) and the year of birth is also desired.

MBL/WHOI LIBRARY



WH 17KP 6

